

The Oligocene succession in the eastern North Sea: basin development and depositional systems

E. M. JARSVE*[†], T. EIDVIN[‡], J. P. NYSTUEN*, J. I. FALEIDE*,
R. H. GABRIELSEN* & B. I. THYBERG[§]

*Department of Geosciences, University of Oslo, PO Box 1047 Blindern, N-0316 Oslo, Norway

[‡]Norwegian Petroleum Directorate, PO Box 600, NO 4003 Stavanger, Norway

[§]CGG Multi Client Scandinavia and Russia, PO Box 490 Skøyen, 0213 Oslo, Norway

(Received 19 February 2014; accepted 10 September 2014)

Abstract – The Oligocene sedimentary succession in the eastern North Sea is revised and re-interpreted by applying new state-of-the-art reflection seismic data integrated with new bio- and Sr-stratigraphy data from three key wells in the study area. The Oligocene succession in the eastern North Sea is divided into four transgressive–regressive (T-R) sequences, characterized by non-accretional and/or aggradational transgressive systems tracts and prograding regressive systems tracts. Detailed studies of three wells, including biostratigraphy and Sr analysis, constrain the age relationships between the T-R sequences. Internal clinof orm geometry indicates that the sediments were sourced from the present southern Norwegian mainland to the north of the depositional area. The direction of progradation shifted from being SE-directed in the earliest Rupelian (early Oligocene) to S- and SW-directed during Chattian time (late Oligocene). Rapid basin subsidence is indicated by the development of non-accretionary transgressive systems tracts, with subsequent progradation into water depths of hundreds of metres. The creation of accommodation space was out of phase relative to eustatic sea-level changes, and mainly controlled by regional-scale differential vertical movements where uplift and exposure of landmasses of the hinterland (southern Norway) occurred concurrently with basin subsidence. Halokinesis had an intra-basinal influence on the main sediment transport direction, but probably did not contribute much in creation of accommodation space.

Keywords: sequence stratigraphy, sedimentology, structural geology, source to sink.

1. Introduction

To understand the source to sink relationships in a sedimentary basin it is crucial to comprehend the coupling between regional and local tectonics, together with climate changes and hinterland geomorphology. Interest has therefore increased on the onshore–offshore relationship to study the relation between climate and tectonics (e.g. Riis, 1996; Gabrielsen *et al.* 2005, 2010; Stoker *et al.* 2005; Japsen *et al.* 2006; Nielsen *et al.* 2009; Medvedev, Souche & Hartz, 2013; Miller *et al.* 2013; Redfield & Osmundsen, 2013).

It is evident from observations of siliciclastic wedges in the eastern North Sea basin that the source area of terrigenous detritus changed from being situated to the west during Paleocene and Eocene times to becoming mainly situated to the north (i.e. southern Norway) during Oligocene time (e.g. Jordt *et al.* 1995; Jordt, Thyberg & Nøttvedt, 2000; Michelsen *et al.* 1995; Faleide *et al.* 2002). It has previously been proposed that observed changes in the sedimentary pattern at the Eocene–Oligocene boundary of North Europe was due to tectonic uplift of Scandinavia and associated subsidence in the adjacent North Sea basin (e.g. Jordt *et al.*

1995; Michelsen *et al.* 1995; Faleide *et al.* 2002; Gabrielsen *et al.* 2010). The same authors suggested that tectonic increase in relief was the triggering mechanism of changes in climatic conditions and associated switches in erosion and drainage systems. In a series of recent studies this view has however been challenged by the interpretation that the recorded changes in sedimentary trends can be ascribed solely to major climatic cooling at the Eocene–Oligocene boundary (e.g. Clausen *et al.* 1999; Huuse & Clausen, 2001; Huuse, Lykke-Andersen & Michelsen, 2001; Nielsen *et al.* 2009, 2010; Gole-dowski, Nielsen & Clausen, 2012). Nielsen *et al.* (2009, 2010) suggest that the topography of southern Norway has been significant since the establishment of the Caledonian Orogeny during Silurian–Devonian time, and mainly changed through the combined effects of erosion due to climatically forced weathering/erosion and associated isostatic response and extensional collapse. This view therefore excludes the effect of increased relief from tectonic uplift of Scandinavia during Oligocene time.

In our view, several aspects of the tectono-dynamic and climatic changes that influenced Cenozoic sedimentation of the eastern North Sea are not yet fully understood, such as the relationship between the creation of accommodation space and global climate changes

[†] Author for correspondence: e.m.jarsve@geo.uio.no




Figure 1. (a) Location of the study area. (b) Structural elements within the study area, including wells applied in this study (CG – Central Graben; DCS – Danish Continental Shelf; CSFC – Coffee–Soil Fault Complex; EB – Egersund Basin; FFZ – Fjerrislev Fault Zone; HG – Horn Graben; KFZ – Kreps Fault Zone; LFB – Lista Fault Blocks; NCS – Norwegian Continental Shelf; NDB – Norwegian Danish Basin; RFH – Ringkøbing Fyn High; SG – Skagerrak Graben; STZ – Sorgenfrei Torquist Zone; ÅG – Åsta Graben). See correlation between Norwegian wells 11/10–1, 9/12–1 and 2/2–2 in Figure 4. (c) Seismic data coverage available for this study (courtesy of Fugro Multi Client Services and TGS Nopec).

and basin subsidence. In this context, the Oligocene succession studied provides relevant and new information on sediment transport directions, sedimentation rates and migration of depocentres, and is a good candidate for studying the relationship between tectonic and climatic influence in a source to sink system. In particular, we emphasize the development of accommodation space in the basinal area and its possible link to the development of relief in the hinterland area and/or climate changes during Oligocene time via integrated studies of bio- and Sr-stratigraphy, seismic sequence stratigraphic techniques and published climate data. The present work therefore aims to re-evaluate the Oligocene sedimentary succession in the eastern North Sea area, based on improved age control and

enhanced seismic coverage in the Norwegian sector (Fig. 1). These data are complementary to the dataset used in previous studies (Michelsen & Danielsen, 1996; Danielsen, Michelsen & Clausen, 1997), which had their main focus on the Oligocene sequences in the Danish sector. Our study includes an analysis of any relative influence of tectonic uplift and climate changes on the development of relief and the rate of erosion in the hinterland and sedimentary flux to the basin. We also discuss the processes influencing the accommodation space in the basin such as basin subsidence, eustasy and halokinesis. Furthermore, we discuss the interplay between climate change and tectonics from seismic sequence stratigraphic analyses. The study area is situated within longitude 3–8° E and latitude 55–58° N (Fig. 1).

2. Regional geology

The main structural element in the study area is the Norwegian–Danish Basin, which is a major NW–SE-oriented basin of Mesozoic age in the central and eastern North Sea. It is bounded to the south by the Ringkøbing–Fyn High and the Coffee–Soil Fault Complex and to the north by the Fjerritslev Fault Zone and Farsund Basin (Fig. 1). The Fjerritslev Fault Zone and Farsund Basin are closely associated with the Sorgenfrei–Tornquist Zone, as known from Denmark and further south-eastwards, and are often considered as the Fennoscandian Border Zone (Ziegler, 1990; Berthelsen, 1998; Fredriksen, Nielsen & Balling, 2001; Heeremans & Faleide, 2004; Heeremans *et al.* 2004).

One of the most striking events in the North Sea basin during Cenozoic time was the marked change in sediment provenance at the Eocene–Oligocene boundary in the eastern North Sea area (e.g. Jordt *et al.* 1995; Jordt, Thyberg & Nøttvedt, 2000; Huuse, 2002; Faleide *et al.* 2002; Anell, Thybo & Stratford, 2010). During latest Paleocene–Eocene time, sediments of the North Sea basin were primarily derived from the East Shetland Platform and the British Isles to the west (Galloway *et al.* 1993; Jordt *et al.* 1995; Rohrman *et al.* 1995; Faleide *et al.* 2002; Anell, Thybo & Stratford, 2010; Anell, Thybo & Rasmussen, 2012). This was related to tectonic uplift of the East Shetland Platform during the late Paleocene break-up of the NE Atlantic (e.g. Jordt *et al.* 1995; Faleide *et al.* 2002). Minor sediment influx from the north during Paleocene time in the eastern North Sea area (e.g. Hamberg *et al.* 2005) may also indicate a tectonic uplift of parts of southern Norway during the break-up of the NE Atlantic (Jordt *et al.* 1995; Jordt, Thyberg & Nøttvedt, 2000; Martinsen *et al.* 1999; Huuse, 2002; Faleide *et al.* 2002; Dmitrieva *et al.* 2012). The main sediment transport was focused towards the eastern North Sea during early Oligocene time, reflecting a shift from a mainly western source area during Paleocene–Eocene time to a mainly northern source area during Oligocene time (e.g. Jordt *et al.* 1995; Michelsen *et al.* 1995; Clausen *et al.* 1999; Jordt, Thyberg & Nøttvedt, 2000; Faleide *et al.* 2002; Japsen, Bidstrup & Lidmar-Bergström, 2002; Gabrielsen *et al.* 2005, 2010; Japsen *et al.* 2007; Schiøler *et al.* 2007; Anell, Thybo & Artimieva, 2009; Anell, Thybo & Stratford, 2010; Anell, Thybo & Rasmussen, 2012). The same authors ascribe this shift in main provenance area to a tectonic uplift of southern Norway during late Palaeogene–Neogene times. A second tectonic phase was proposed to have occurred during middle Miocene time (e.g. Japsen & Chalmers, 2000; Stoker *et al.* 2005; Japsen *et al.* 2007; Rasmussen, 2009). However, other authors have suggested that the Oligocene siliciclastic wedges and middle Miocene unconformity are related to climatic changes, reflected by increased erosion and sediment transport (e.g. Huuse, Lykke-Andersen & Michelsen, 2001; Huuse, 2002; Goleadowski, Nielsen & Clausen, 2012). These authors suggest that only isostasy, climate and eustatic

changes influenced erosion and sediment transport on an already-established topography. Nielsen *et al.* (2009, 2010) also believe that the then existing topography in southern Norway was a remnant of the Caledonian Orogeny (the isostasy–climate–eustasy or ICE hypothesis).

During Pliocene–Pleistocene time, glaciations in Scandinavia gave rise to increased erosion rates of the Norwegian mainland, with subsequent isostatic response and basinward tilting (i.e. dip-direction to the SW) of the Mesozoic and Cenozoic sedimentary successions in the eastern North Sea (e.g. Riis, 1996; Huuse, Lykke-Andersen & Michelsen, 2001; Faleide *et al.* 2002; Huuse, 2002; Gabrielsen *et al.* 2005, 2010; Nielsen *et al.* 2009). However, Stoker *et al.* (2005) showed that the onset of major progradation was initiated at about 4 Ma, which predates the growth of glaciation by at least 1 Ma, implying a potential tectonic phase during Pliocene time (Japsen *et al.* 2007).

3. Seismic sequence stratigraphy

3.a. Data

The present study utilizes more than 20 000 km of long-offset 2D data (North Sea Renaissance survey), combined with detailed studies of three key wells (11/10–1, 9/12–1 and 2/2–2) in the Norwegian sector (Fig. 1). The NSR seismic survey was acquired by Fugro and TGS Nopec during the period 2003–2009 on the Norwegian continental shelf. The NSR seismic survey is a high-quality uniform dataset with a vertical resolution of *c.* 10–20 m, which offers seismic coverage beyond that seen in previous studies. In addition to the NSR survey, the NDT and UG surveys in the Norwegian sector and DCS, DK1, DK2, DT97, SP82 and UCGE97 surveys in the Danish sector were utilized to achieve a complete updated coverage of the study area and to perform a full comparison with previous studies.

3.b. Previously applied seismic stratigraphic sequences

The Cenozoic succession in the northern and eastern North Sea area has been subdivided into depositional sequences *sensu* Mitchum, Vail & Thompson (1977) and Vail (1987) (Exxon Model) by Jordt *et al.* (1995), Michelsen & Danielsen (1996), Danielsen, Michelsen & Clausen (1997), Michelsen *et al.* (1998), Martinsen *et al.* (1999), Jordt, Thyberg & Nøttvedt (2000) and Faleide *et al.* (2002). However, these authors pointed out the challenges in identifying the system tracts that ought to be present according to the depositional sequence model of Exxon, particularly the lowstand and transgressive systems tracts (Michelsen *et al.* 1995; Michelsen & Danielsen, 1996; Danielsen, Michelsen & Clausen, 1997; Michelsen *et al.* 1998; Martinsen *et al.* 1999). Michelsen & Danielsen (1996) and Danielsen, Michelsen & Clausen (1997) documented that downlap surfaces correlated with high gamma peaks and interpreted these surfaces as maximum flooding surfaces. However, they were also able to identify the sequence boundary (*sensu* Exxon depositional model)




Figure 2. (Colour online) Sketch summarizing the four T-R sequences described in this study and the corresponding sequence boundaries (SB), transgressive surfaces (TS) and maximum flooding surfaces (MFS). Note the non-accretionary nature of the TST of OSS-1 and OSS-4, while the TST in OSS-2 and OSS-3 are aggradational. The sketch is not to scale.

on the basis of both seismic downlaps and onlaps, which appears contradictory as downlap presumes the presence of accommodation space above the surface. If the boundaries of the sequences defined by Michelsen *et al.* (1995, 1998) (and in the other studies referred to above that followed the same systematics) were formed as subaerial unconformities in accordance with the Exxon depositional sequence model (*sensu* Van Wagoner *et al.* 1987, 1988, 1990; Posamentier, Jervey & Vail, 1988; Posamentier & Vail, 1988), there would not be any marine accommodation space above these surfaces except in the deeper parts of the basins, and then above conformable surfaces correlative to the subaerial unconformity. Nevertheless, above some of these sequence boundaries there are downlapping prograding successions, implying that marine accommodation had formed.

3.c. Seismic sequence stratigraphic model applied in the present study

The challenges related to the identification of bounding surfaces between possible lowstand and transgressive deposits on seismic sections in Cenozoic successions in the North Sea area make it difficult to apply the lowstand, transgressive and highstand systems tracts of the Exxon depositional sequence model. However, as also pointed out by Michelsen *et al.* (1998), several of the Cenozoic successions display distinct transgressive–regressive (T-R) trends, demonstrated by seismic facies, clinoform geometry and well-log trends. Transgressive–regressive trends reflect the interplay between the creation and/or destruction of accommodation space and the rate of sedimentation, the controlling factors in basin infill dynamics irrespective of types of allogenic driving factors such as tectonics, eustasy, climate and geomorphology. In our sequence stratigraphic analysis we therefore apply the T-R sequence stratigraphic model (Embry, 1988, 1993, 1995; Embry & Johannessen, 1992; Fig. 2).

The transgressive systems tract (TST) can be non-accretionary, implying a lack of sediments deposited during the rise in relative sea level, or consisting of

sediment thicknesses less than the seismic resolution (Helland-Hansen, 1995). Accretionary TST consists of marine successions with aggrading stacking geometry (Fig. 2). The underlying bounding transgressive surface (TS) coincides with a regional unconformity, defined by toplap and/or toplap truncation. The part of the surface characterized by toplap truncation is interpreted as a subaerial unconformity (SU), whereas the toplap part of the surface is a candidate for the marine conformable part of the SU. Alternatively, this part of the regional unconformity may represent subaerial exposure without marked erosion or a shallow marine omission, bypass or ravinement surface. Onlap on the regional unconformity implies creation of accommodation space and hence the establishment of the TS, accompanied by sedimentation, as marine and/or coastal onlap. The combined subaerial unconformity and the transgressive surface are termed a SU/TS surface.

The transgressive systems tract is bounded at the top by a regional surface characterized by downlap. This is the maximum flooding surface (MFS), forming the lower bounding surface of the regressive systems tract (RST). In cases where a transgressive systems tract is absent, the MFS may coincide with both the regional unconformity and the transgressive surface, forming a combined SU/TS/MFS surface. The regressive systems tracts are distinguished by progradational successions with internal clinoform geometry. In this system, eventual deposits formed during a fall in relative sea level will be included in the regressive systems tract.

3.d. Nomenclature

In this study, we introduce an adjusted nomenclature for the Oligocene seismic sequences (OSS; Figs 2, 3) relative to previous studies (Jordt *et al.* 1995; Michelsen & Danielsen, 1996; Danielsen, Michelsen & Clausen, 1997). New nomenclature is needed because the T-R systematic subdivision is employed in this study, whereas previous studies applied the Exxon systematics. We divide the Oligocene succession into four seismic sequences, where two of the sequences are of early Oligocene age (OSS-1 and OSS-2) and the two later




Figure 3. (Colour online) Seismic stratigraphic sequences correlation with the NSA zonation of King (1989) and Gradstein & Bäckström (1996), correlated to that by Michelsen & Danielsen (1996) and Jordt, Thyberg & Nøttvedt (2000). Arrows indicate direction of sediment progradation.

sequences are of late Oligocene age (OSS-3 and OSS-4). The sequence boundaries are compared to previous studies and summarized in Figure 3.

4. Methods

4.a. Well correlation

The seismic data were tied to available wells, including key wells 11/10-1, 9/12-1 and 2/2-2 in the Norwegian sector (Figs 4, 5) and wells F-1, Inez-1, D-1, Nini-1 and Ibenholt-1 in the Danish sector. Extensive studies, including biostratigraphy and Sr-stratigraphy, were carried out on material from the Norwegian wells. More comprehensive studies of the Norwegian wells are presented in Eidvin *et al.* (2013). In combination with new information obtained from the biostratigraphical and chemostratigraphical studies, the seismic geometries were used to establish a depositional model for the study area. A particular emphasis was placed on recognizing progradation direction and transgressions, as well as maximum flooding surfaces.

4.b. Biostratigraphy and strontium isotope stratigraphy (SIS)

For this study, new micropalaeontological and Sr-isotope analyses of well 2/2-2, 9/12-1 and 11/10-1 are used to obtain better age control on the Oligocene se-

quence in the eastern North Sea. A total of 66 ditch cutting samples and two conventional core samples (well 2/2-2) were analysed (Table 1, Fig. 4).

4.b.1. Micropalaeontological analyses

The micropalaeontological investigations are based on analyses of planktonic and benthic foraminifera and pyritized diatoms. The fossil assemblages are correlated with the biozonation of King (1989), who outlines a micropalaeontological zonation for Cenozoic sediments in the North Sea. Gradstein & Bäckström's (1996) faunal zonation of the North Sea and Haltenbanken is also used.

4.b.2. Strontium isotope analyses

Strontium isotope stratigraphy is used as an additional control on the biostratigraphic correlations. The $^{87}\text{Sr}/^{86}\text{Sr}$ ratio of seawater is very uniform on a global scale, which is a reflection of the long oceanic residence time of strontium (2–4 Ma) combined with a relative short (≤ 2000 a) oceanic mixing rate. Strontium isotope stratigraphy is an effective tool, particularly for the dating of Miocene and Oligocene sections. It has best resolution in sediments older than 15 Ma because the Sr isotopic composition of seawater changed rapidly during this period (e.g. Howarth & McArthur 1997).

A total of 23 samples were investigated for their Sr isotopic compositions with 41 analyses (Table 1, Fig. 4). The majority (26 analyses) were conducted on tests of calcareous index foraminifera and the remainder on mollusc fragments (15 analyses from sandy sections). Strontium values were converted to age estimates using the SIS look-up table of Howarth & McArthur (1997). This table is based on the timescale compiled by Berggren *et al.* (1995), which for the Oligocene does not deviate significantly from the new timescale of the International Commission on Stratigraphy (see <http://www.stratigraphy.org/ICSChart/ChronostratChart2013-01.pdf>). For more details on the precision of the method, see Howarth & McArthur (1997), McArthur, Howarth & Bailey (2001) and Eidvin & Rundberg (2001, 2007). Note that the micropalaeontological zonation of King (1989) is based on the timescale of Berggren, Kent & Van Couvering (1985), but we have converted the ages to the timescale of Berggren *et al.* (1995). The micropalaeontological zonation of Gradstein & Bäckström (1996) is based on the timescale of Cande & Kent (1992) in which the absolute ages are identical to those of Berggren *et al.* (1995).

4.c. Volume estimates and water depths

The sound wave velocity is set to 2000 m s^{-1} for the combined Cenozoic succession, as measured in the studied wells (check shot; 11/10-1, 9/12-1 and 2/2-2). A velocity of 2000 m s^{-1} was also applied by Jordt,

Table 1. Strontium isotope data from well 11/10–1, 9/12–1 and 2/2–2, analysed at the University of Bergen. Sr ratios were corrected to NIST 987 = 0.710248. The numerical ages were derived from the SIS look-up table version 3:10/99 of Howarth and McArthur (1997). NIST – National Institute for Standards and Technology.

Lithostratigraphic unit	Sample (ditch cuttings, m)	Corrected $^{87}\text{Sr}/^{86}\text{Sr}$	2σ error	Age (Ma)	Comments	Analysed fossils
Well 11/10–1						
Vade Formation	304.8	0.708177	0.000007	25.26		9 large tests of <i>Elphidium subnodosum</i>
Vade Formation	304.8	0.708118	0.000008	26.48		10 large tests of <i>E. subnodosum</i> , <i>Astigerina guerichi guerichi</i> and <i>Guttulina</i> sp.
Vade Formation	365.8	0.708091	0.000008	27.23		1 mollusc fragment
Vade Formation	365.8	0.708101	0.000007	26.93		1 mollusc fragment
Vade Formation	426.7	0.708008	0.000009	29.39		1 mollusc fragment
Vade Formation	426.7	0.708227	0.000009	24.48	Caved?	1 mollusc fragment
Vade Formation	457.2	0.707979	0.000008	30.11		Approximately 30 fragments of <i>Bryozoa</i> and <i>Echinoderma</i>
Vade Formation	472.4	0.708039	0.000012	28.61		50 small tests of <i>Pararotalia canui</i> , <i>G. angustumbilicata</i> and <i>G. Praebulloides</i>
Hordaland Group	487.7	0.708015	0.000008	29.22		1 mollusc fragment
Hordaland Group	487.7	0.707991	0.000007	29.81		1 mollusc fragment
Hordaland Group	502.9	0.707991	0.000008	29.81		1 mollusc fragment
Hordaland Group	502.9	0.707973	0.000007	30.27		1 mollusc fragment
Hordaland Group	518.2	0.707951	0.000008	30.87		1 mollusc fragment
Hordaland Group	518.2	0.707937	0.000008	31.24		1 mollusc fragment
Hordaland Group	640.1	0.707938	0.000007	31.22		9 large tests of <i>G. soldanii girardana</i> , <i>Nodosaria</i> sp. A (King, 1989), <i>Lagena</i> sp., <i>Alabama scitula</i>
Hordaland Group	640.1	0.708077	0.000008	27.69	Caved?	1 mollusc fragment
Hordaland Group	640.1	0.708250	0.000008	24.10	Caved?	1 mollusc fragment
Hordaland Group	823.0	0.707875	0.000008	32.59		19 tests of <i>Stilostomella</i> spp., <i>T. alsatica</i> , <i>R. bulimoides</i> , <i>Ceratobulimina contraria</i> , <i>G. soldanii girardana</i> , <i>Cibicides mexicanus</i>
Hordaland Group	853.4–883.9	0.707871	0.000008	32.66		26 tests of <i>Stilostomella</i> spp., <i>Bolivina</i> cf. <i>antique</i> , <i>T. alsatica</i> , <i>R. bulimoides</i> , <i>Hoeglundina elegans</i> , <i>G. soldanii girardana</i> , <i>Cibicides mexicanus</i>
Hordaland Group	914.4	0.707871	0.000008	32.66		Twelve tests of <i>G. soldanii mamillata</i> , <i>H. elegans</i> , <i>G. soldanii girardana</i> , <i>Stilostomella</i> spp., <i>G. subglobosa</i> , <i>Fronicularia budensis</i> , <i>T. alsatica</i> , <i>C. contraria</i> , <i>G. praebulloides</i>
Well 9/12–1						
Hordaland Group	463.3	0.708135	0.000008	26.10		1 mollusc fragment
Hordaland Group	463.3	0.708150	0.000008	25.79		1 mollusc fragment
Hordaland Group	475.5	0.708152	0.000008	25.75		1 mollusc fragment
Hordaland Group	475.5	0.708145	0.000008	25.89		1 mollusc fragment
Hordaland Group	475.5	0.708140	0.000007	25.99		1 mollusc fragment
Hordaland Group	487.6	0.708103	0.000008	26.87		1 mollusc fragment
Hordaland Group	487.6	0.708159	0.000008	25.61		Approximately 20 tests of <i>A. guerichi guerichi</i> , <i>A. guerichi staeschei</i>
Hordaland Group	499.9	0.708080	0.000008	27.55		1 mollusc fragment
Hordaland Group	499.9	0.708128	0.000009	26.25		1 mollusc fragment
Hordaland Group	499.9	0.708111	0.000008	26.65		1 mollusc fragment
Hordaland Group	512.1	0.708231	0.000008	24.42		Approximately 15 tests of <i>A. guerichi guerichi</i> , <i>A. guerichi staeschei</i>
Hordaland Group	512.1	0.708136	0.000008	26.08		1 mollusc fragment
Hordaland Group	512.1	0.708113	0.000007	26.60		1 mollusc fragment
Hordaland Group	512.1	0.708114	0.000008	26.57		1 mollusc fragment
Hordaland Group	524.3	0.708092	0.000008	27.20		Approximately 35 tests of <i>A. guerichi guerichi</i> , <i>A. guerichi staeschei</i>
Hordaland Group	536.5	0.708075	0.000008	27.69		Approximately 23 tests of <i>A. guerichi guerichi</i> , <i>A. guerichi staeschei</i>
Hordaland Group	548.6	0.708103	0.000007	26.87		Approximately 55 tests of <i>A. guerichi guerichi</i> , <i>A. guerichi staeschei</i>
Hordaland Group	573.0	0.708042	0.000007	28.53		Approximately 40 tests of <i>A. guerichi guerichi</i> , <i>A. guerichi staeschei</i>
Hordaland Group	609.6	0.708037	0.000007	28.66	Reworked?	10 tests of <i>Siphonina reticulata</i>
Well 2/2–2						
Vade Formation	1984.5	0.708127	0.000009	26.3		Fragment of 1 large bivalve
Vade Formation	1984.5	0.708127	0.000009	26.0		Fragment of 1 large bivalve




Figure 4. Correlation between the three wells on the Norwegian Continental Shelf applied to this study for Sr-stratigraphy and biostratigraphic analysis.

Thyberg & Nøttvedt (2000). Volumes of sediments are calculated for each of the sequences within the 100 m (100 ms two-way travel time or TWT) contours (Fig. 6). The volumes are calculated between the upper and lower bounding surfaces for each sequence, hence including both the TST and the RST. Note however that these estimates give minimum volumes, because no allowance has been made for compaction and

post-depositional erosion. In addition, there may be undetected sediment volumes deposited beyond the study area.

The estimated water depths are based on the heights of clinoforms for each sequence as calculated after correcting for later tilting of the corresponding downlap surface, assuming that the clinoform height also represents the palaeowater depth.




Figure 5. Seismic correlation between the three key wells 11/10–1, 9/12–1 and 2/2–2 in this study.

5. Description and analysis of the seismic sequences

The Oligocene seismic succession was subdivided into four second-order seismic sequences, all of which are characterized as T-R sequences (Fig. 2). The following sections describe the four Oligocene seismic sequences (OSSs).

5.a. Oligocene seismic sequence 1 (OSS-1)

The lower boundary of OSS-1 is defined by a downlap surface (Fig. 7). This surface is characterized by a strong and continuous reflector which can be mapped across the entire study area. It represents sedimentary strata which drape a surface that shows erosional truncation into underlying Eocene and Paleocene strata to the east and southeast of Norwegian well 9/12–1 (Fig. 8). OSS-1 is identified mainly in the Norwegian sector and is upward bounded by toplap truncation and succeeded by OSS-2 (Fig. 7). The upper sequence unconformity corresponds with the Oi1a global cooling event (Pekar *et al.* 2002), which in well 11/10–1 also corresponds to a section including the dinoflagella cyst *Svalbardella* (Sliwinska and Heilmann-Clausen, 2011).

The main depocentre of OSS-1 is located in the area west and southwest of Norwegian well-site 11/10–1 (Fig. 7). Here the sequence reaches a maximum thickness of 300 m (Fig. 7) and the estimated net depositional volume rate is $866 \text{ km}^3 \text{ Ma}^{-1}$. This is a minimum estimate, as volume loss from mechanical compaction and sediments transported beyond the study area are not included. The sequence thins eastwards to less than 50 m at the Danish well F-1, and also thins towards the west and southwest (Fig. 7).

In the central part of the depocentre, OSS-1 is strongly influenced by a salt structure (Fig. 9). This corresponds to the area where the sequence reveals its maximum thickness. OSS-1 strata thin across this salt structure and onlap its flanks (Fig. 9). Similarly,

the sequence onlaps a salt structure southeast of well 11/10–1 on the Norwegian–Danish border, and further to the east its thickness is reduced to less than that detectable within the seismic resolution. OSS-1 is not seen in the Inez-1 well area. Towards the north, OSS-1 onlaps Eocene strata along the NW–SE-striking Fjerritslev Fault Zone (Figs 10, 11a). Note that, in contrast to OSS-1, the Eocene and Paleocene sequences thicken northwards, indicating that the Fjerritslev Fault Zone area represented parts of the depositional area during early Palaeogene time.

OSS-1 contains seismic geometries that are characterized by clinoforms prograding SSE, which is perpendicular to the basin axis during deposition (NE–SW; Fig. 7). The clinoforms have a complex sigmoidal/oblique geometry, with dips of $0.5\text{--}1^\circ$. The clinoform heights are approximately 100 m, and the prograding succession is upwards bounded by clinoform toplap truncation (SU). The clinoforms shift to a more sigmoidal geometry in the southwest part of the depositional area, where the clinoform topsets are preserved (Fig. 11b). Well 11/10–1, which penetrates the prograding clinoforms, reveals gamma log readings consistent with fine-grained sediments throughout OSS-1 (Fig. 4). The gamma log also reveals an upwards-decreasing gamma ray response of individual packages, interpreted as upwards-coarsening parasequences, typical of prograding regressive depositional systems.

5.a.1. Interpretation

Based on the downlapping of the internal clinoforms onto the basal bounding reflector of OSS-1 and the strong and regional character, this basal reflector is interpreted as a MFS. The Eocene–Oligocene boundary at the base of OSS-1 resembles a marine erosional surface, representing either a ravinement surface (>20 m water depth) or the action of deep marine currents (up to hundreds of metres water depth, e.g. Nichols, 2009).




Figure 6. (Colour online) Seismic stratigraphic subdivision from this study, with characteristic features for each sequence. Note that water depth is calculated from clinoform heights.

However, as the TST at the base of OSS-1 is non-accretionary, the lower bounding surface of OSS-1 is a combined SU/TS/MFS surface, at least in the platform margin areas of the basin. This is supported by well data offshore Denmark (Danielsen, Michelsen & Clausen, 1997) and by observations onshore Denmark. These reveal that the base of the Viborg Formation contains glauconite, supporting the presence of a well-developed hiatus (Christensen & Ulleberg, 1973) that may represent a non-accretionary TST.

The clinoform heights indicate water depths of approximately 100 m during earliest Rupelian time. The shale lithology, as recorded in well 11/10-1, is consistent with a parallel and sigmoidal clinoform geometry together with gentle clinoform dips (0.5–1°), typical where fine-grained material dominates (e.g. Mitchum,

Vail & Thompson, 1977; Veeken, 2007). Such depositional systems are also often characterized by low energy in the basin, where limited reworking of the sediments takes place.

The thinning of OSS-1 and its onlapping pattern against the salt diapir located within the central part of the OSS-1 depocentre (Fig. 9) is taken to indicate diapirism during deposition of the sequence during early Rupelian time. Similarly, halokinesis is interpreted to have controlled the eastern extent of the main depositional area of OSS-1. This is seen from the onlap contact and thinning of OSS-1 onto salt diapirs at the Norwegian–Danish border, reflecting bathymetric relief here during earliest Rupelian time. Salt mobilization is also known from onshore Denmark during Oligocene time (Japsen *et al.* 2007).




Figure 7. (Colour online) (a) Time-thickness map of OSS-1. The clinoform facies for the various units represents the highstand systems tract and also the main depocentre at the time of deposition. (b) The upper surface of OSS-1 is marked in red in the seismic section (B–B'). (c) Seismic termination above offlap break, indicating clinoform toplap truncation.

5.b. Oligocene seismic sequence 2 (OSS-2)

The lower boundary of OSS-2 corresponds to the erosional surface representing the SU at the top of OSS-1. OSS-2 also contains an internal downlap surface, represented by a strong continuous seismic reflector which is recorded across most of the study area (Fig. 7). The upper boundary of OSS-2 is characterized by toplap truncation and severe incision (Fig. 12), and

is considered equivalent to the upper boundary of unit 4.2 in Michelsen & Danielsen (1996; Fig. 3).

OSS-2 displays a maximum thickness of approximately 550 m south of wells 11/10–1 and F-1, and thins south- and westwards to less than 100 m (Fig. 12). Unlike OSS-1, OSS-2 maintains a uniform thickness northwards with no observed onlap or marked thinning in the Fjerritslev Fault Zone area (Fig. 10). Based on the




Figure 8. (Colour online) Seismic terminations indicating erosion of Eocene strata at base OSS-1. Note the strong seismic amplitude directly above the unconformity, corresponding to the downlap surface of OSS-1.

recorded thicknesses, the bulk sediment volume rate is calculated as *c.* 616 km³ Ma⁻¹ (Fig. 6).

In well 11/10–1 and southwards, the lower part of OSS-2 is characterized by seismic geometries showing SE-dipping clinoforms with offlap breaks defining a near-vertical trajectory, showing an aggrading depositional geometry of *c.* 100 m (Fig. 10). The gamma ray log trend in well 11/10–1 indicates the presence of slightly upwards-coarsening parasequences (Fig. 4). The gamma response culminates in the middle part of OSS-2 in a strong peak (Fig. 4) which, in stratigraphic position, coincides with a strong and continuous seismic reflection (Fig. 5). This continuous character of the seismic reflection likely reflects strata that drape the aggrading clinoforms and acts as a downlap surface for the upper part of OSS-2.

The upper part of OSS-2 is characterized by a seismic succession with prograding clinoforms dipping towards the south and southwest. The clinoforms have a tangential oblique geometry, with clinoform heights of approximately 200–250 m and clinoform dips of 1.5–5.5°. The upper boundary suggests erosion as evidenced by truncation and incision of the clinoform topsets (Fig. 12). The incision features on the clinoform topsets have vertical depths of up to 20–30 m (Fig. 12), indicating channel deposition. The clinoforms shift westwards to more complex sigmoidal/oblique geometry (Fig. 11b). Onlap from sediments belonging to OSS-3 is recorded against the upper SU of OSS-2, indicating coastal onlap (Fig. 12).

Internally, the clinoform system in the upper part of OSS-2 is highly complex with lensoidal subunits bounded by unconformities and onlap surfaces (Fig. 12), indicating internal clinoform reflectors dipping in various directions.

In the frontal part of the prograding clinoforms (i.e. basinwards), OSS-2 is characterized by a 150 m interval with strong seismic amplitude reflectors (Figs 10, 11a). These seismic reflectors downlap onto the bottom-bounding reflector relative to the succession of SSW-prograding clinoforms of the same sequence, as well as onlapping the frontal part of the prograding clinoforms towards the NNE (Fig. 10).

At the position of Norwegian wells 11/10–1 and 9/12–1 and northwards, OSS-2 is characterized by par-

allel seismic reflectors. The gamma ray log in well 11/10–1 reveals an upwards-coarsening succession, as indicated by the upwards decrease in gamma readings (Fig. 4).

5.b.1. Interpretation

The lower bounding surface for OSS-2 represents a sequence boundary separating OSS-1 and OSS-2. As indicated by the aggrading depositional pattern, the lowermost part of OSS-2 is interpreted as a TST where sediment input closely balanced the rate of creation of accommodation space during the transgression. The strong and continuous character of the reflector, its downlap pattern and the gamma ray peak of the reflector defining the top of the TST are all interpreted to represent the MFS. As indicated by the clinoform heights, the water depth during transgression is assumed to have reached approximately 250 m during maximum transgression.

The prograding part of OSS-2 is interpreted to constitute the RST. The oblique geometry of the clinoforms is typical of coarse-grained clastic debris (Hansen & Rasmussen, 2008; Bullimore *et al.* 2005). However, the internal architecture of the upper part of OSS-2 with its shifting lensoid geometry and erosional surfaces (Fig. 12) probably represents various depositional lobes, shifting in position and progradational direction through time. Lateral shifts in sediment accumulation are typical of delta lobes and can be explained by avulsion and shifting transport directions of the fluvial system in the hinterland, as seen in modern deltas (e.g. the Mississippi delta; Coleman, 1988). However, as halokinesis is recorded at the time of deposition of OSS-2, local doming within the basin is likely to have created local barriers for the sediments and therefore also influenced the pattern of sedimentation laterally through time.

The sediments belonging to the TST of OSS-2 entered the basin from the NNW, following similar sediment routing as OSS-1. The shift in direction of progradation, from southeast for OSS-1 and the lower part of OSS-2 to south and southwest for the upper part of OSS-2, may indicate that the entry point of the sediment influx to the basin shifted or that depocentres




Figure 9. (Colour online) (a) Base map showing salt structures in the study area penetrating and/or affecting the Oligocene sequences. Seismic sections showing halokinesis during deposition of (b) OSS-1 and (c) OSS-3. Arrows mark seismic onlap against the salt structures.

of depositional lobes changed position. The parallel subhorizontal seismic reflectors in the northern part of the study area are interpreted as shallow marine shelf deposits.

Based on the downlap onto the maximum flooding surface and onlap onto the RST of OSS-2, the up-

permost part of OSS-2 is interpreted as lowstand fan deposits. These are likely to contain gravity flow facies derived as erosional material from the shelf (Posamentier & Vail, 1988).

The depositional evolution of OSS-2 was terminated by toplap truncation and major incision, creating




Figure 10. Arbitrary seismic sections (striking NNE–SSW) illustrating the various systems tracts of the Oligocene sequences. Note the onlap pattern of OSS-1 against Eocene strata in the NE part. See Figure 1 for location of the seismic lines.




Figure 11. (Colour online) Seismic examples including well tie to well 11/10–1 and the prograding RSTs in the western part of the main depocentre relative to sequences (a) OSS-1 and (b) OSS-2. See Figure 7 for line locations. Note that OSS-1 onlap Eocene strata towards the north, whereas OSS-2 has a more uniform thickness northwards.

an erosional surface interpreted as a sequence unconformity and a sequence boundary (Fig. 12). The upper sequence unconformity is probably of subaerial or shallow marine origin, which is supported by fossil assemblages typical of the upper shoreface in well 11/10–1 (Fig. 4), and potential fluvial incision (Fig. 12).

5.c. Oligocene seismic sequence 3 (OSS-3)

South of the 11/9–1 well area, the lower boundary of OSS-3 bears the characteristics of a marine flooding surface; in the area just south of well 11/10–1, this coincides with the erosional SU bounding OSS-2. Here, downlap of seismic reflectors is recorded in the distal part of the depocentre and the same reflectors onlap the RST of OSS-2 in the more proximal part of the

depocentre (Fig. 10). The upper boundary of OSS-3 is also characterized by toplap truncation (Fig. 13), and corresponds to the top of unit 4.3 in Michelsen & Danielsen (1996; Fig. 3). The main depocentre of OSS-3 was located south of the depocentre of OSS-2, where a maximum thickness of *c.* 200 m (Fig. 13) is recorded. In the southern parts of the study area the lower part of OSS-3 thins across salt diapirs and onlaps against them (Fig. 9b). The thickness and duration of sedimentation are consistent with a sediment volume rate of *c.* 796 km³ Ma⁻¹. The sequence thins to less than 100 m to the south and west, including the 9/12–1 well area (Fig. 13).

In the lower part of OSS-3, seismic reflectors with an aggradational geometry dip towards the SSW. These




Figure 12. (Colour online) (a) Time-thickness map of OSS-2. (b) Clinoform facies for the sequence represents the highstand systems tract and is coherent with the main depocentre at the time of deposition. (c) Internal erosional surfaces in the seismic section and incision at the upper SU.

onlap the prograding clinoforms related to OSS-2, including the erosional surface (SU) which towards the north truncates the OSS-2 clinoform succession (Figs 10, 13).

A strong continuous reflector, which extends across the entire study area, represents sedimentary draping of the aggrading clinoform set. This reflector represents a stratigraphic surface with prograding clinoforms downlapping onto it (Fig. 13). The prograding clinoforms above this surface dip towards the SSW with an oblique geometry, with clinoform toplap truncation below of-flap break (Fig. 13). These prograding clinoforms are restricted to the area of maximum thickness of the sequence (>200 m; Fig. 13) and hence only identified in the Danish sector NNE of the Nini-1 well (Fig. 13). The clinoform height is approximately 100 m, with clinoform dips of 2–3°. Outside this area, the clinoforms shift to a more aggradational style. This is accompanied by a gradual thinning of the seismic successions

to below seismic resolution west and southwest of the Nini-1 well, where the sequence boundary combines with the MFS within the T-R sequence.

In the northern part of the study area at well 11/10–1 and northwards, sequence OSS-3 consists of parallel subhorizontal reflectors. OSS-3 consists mostly of sandstone, with abundant fragments of limestone and molluscs (Fig. 4). Sandstone also dominates the lithology of this sequence in the Danish wells F-1, Inez-1, Ibenholt-1 and D-1 (Danielsen, Michelsen & Clausen, 1997). At the Norwegian well sites 2/2–2 and 9/12–1, which are positioned distally to the main depocentre, OSS-3 consists of shale (Fig. 4).

5.c.1. Interpretation

The upper sequence unconformity of OSS-2 was transgressed and the TS overlain by strata with an internal aggradational geometry of clinoforms. This




Figure 13. (Colour online) (a) Time-thickness map of OSS-3. (b) Clinoform facies for the sequence represents the highstand systems tract and also the main depocentre at the time of deposition. (c) The TST and RST are separated by a MFS.

aggradational succession is interpreted as representing the TST, in which the rate of sedimentation mostly equalled creation of accommodation space (in the same way as for the lower part of OSS-2). The lower boundary of OSS-3 reflects first a shift from lowering of relative sea level with subaerial erosion to stillstand or sediment bypass, succeeded by marine flooding and rise in relative sea level. The southwest dip of the clinoforms indicates that the main progradational direction is from the northeast, perpendicular to the orientation of the basin axis (Fig. 13).

As OSS-3 thins across salt structures in the southern parts of the depositional area, halokinesis appears to have influenced the depositional pattern and acted as barriers in various parts of the basin during deposition of the sequence.

The progradational succession above the MFS within OSS-3 is interpreted to be the RST of OSS-3. The oblique geometry, in addition to the steepness of the clinoforms, is typical of systems deposited in a high-

energy environment where the coarser material may be trapped in the foresets (Bullimore *et al.* 2005; Veeken, 2007). The thickness map and the direction of clinoform dips may indicate that the sequence was sourced from the NNE (Fig. 13). The salt structures, which developed simultaneously as the deposition of OSS-3, likely influenced the continued south-westwards progradation of the succeeding RST by creating a narrow depositional basin for the prograding RST. This is supported by the oblique clinoform geometry, which is often characteristic of narrow depocentres (cf. Bullimore *et al.* 2005). The upper erosional surface that terminates the prograding clinoforms most probably formed as a subaerial unconformity (SU).

5.d. Oligocene seismic sequence 4 (OSS-4)

The lower boundary of OSS-4 is characterized as a downlap surface, where reflectors downlap onto a continuous high-amplitude reflector which extends across




Figure 14. (Colour online) (a) Time-thickness map of OSS-4. (b, c) Internal onlap surfaces, indicating lobe shifting during deposition of the RST of the sequence.

the study area. In the eastern parts of the depocentre, onlap patterns are recorded against the frontal parts of the RST of OSS-3 (Fig. 10). However, this is not observed elsewhere in the depocentre. The clinoforms which onlap the frontal parts of the RST of OSS-3 share the same downlap surface as the overall OSS-4. The upper boundary of OSS-4 is a seismic surface characterized by toplap truncation (Fig. 14), and correlates with the upper boundary of unit 4.4 in Michelsen & Danielsen (1996) and sequence CSS-4 in Jordt *et al.* (1995). OSS-4 has a maximum thickness of 500 m and a sediment volume rate of c. 2546 km³ Ma⁻¹.

The main depocentre of OSS-4 is situated in the Norwegian–Danish Basin and is oriented NW–SE, as seen in the thickness map (Fig. 14). The sequence thins to less than 100 m to the south, southwest and west.

In OSS-4, seismic clinoforms dip towards the southwest and indicate progradation in this direction. The

seismic geometry in the lower part of the sequence shifts from progradational to aggradational eastwards, although with the same direction of dip.

The prograding clinoforms have a sigmoidal geometry, with internal onlap surfaces and erosional surfaces recorded within the succession. However, these are less pronounced and the clinoforms are downlapping the same downlap surface as the overall prograding succession (Fig. 14). The clinoform heights of sequence OSS-4 are 200–300 m, with clinoform dips of 1–2°.

5.d.1. Interpretation

Based on the downlap pattern of the lowermost surface of OSS-4, this is interpreted as a maximum flooding surface (MFS) which, within the seismic resolution, coincides with a transgressive surface (TS). The TS

succeeds a likely subaerial unconformity reflected by fossil assemblages in well 11/10–1, indicating a shallow marine environment, and the toplap truncation that terminates the underlying OSS-3 sequence in the platform areas of the basin. Although the onlap pattern at the frontal part of OSS-3 may represent a thin TST in the eastern part of the study area, the clinofolds share the same downlap surface as the overall sequence. This suggests that the sediment input may have been focused at another part of the basin during progradation. This is supported by the internal onlap surfaces recorded throughout the sequence, which indicate lobe switching and hence changes in direction of main sediment input.

As the transgressive systems tracts is only identified by the MFS at the base of OSS-4 (non-accretionary systems tract), and the maximum flooding surface appears to coincide with the toplap truncation surface cutting the underlying OSS-3, the transgression and resulting increase in accommodation space must have taken place quickly.

The prograding succession above this maximum flooding surface is interpreted to be the RST in OSS-4. The sigmoidal geometry, in addition to the steepness of the clinofolds, is typical of muddy systems, with sediments deposited in a low-energy system with substantial accommodation space (Mitchum, Vail & Thompson, 1977; Bullimore *et al.* 2005). The thickness map and the direction of clinofold dips indicate that the sequence was deposited into the basin from the northeast (Fig. 14).

6. Discussion

The Cenozoic succession in the North Sea area has been the subject of numerous publications in recent years. These publications have mainly been concerned with relating late Palaeogene and Neogene sedimentation variations and depocentre migration to either a tectonic model or climatic model. The tectonic model suggests tectonic uplift of southern Norway (e.g. Jordt *et al.* 1995; Michelsen & Danielsen, 1996; Danielsen, Michelsen & Clausen, 1997; Michelsen *et al.* 1998; Jordt, Thyberg & Nøttvedt, 2000; Faleide *et al.* 2002; Anell, Thybo & Stratford, 2010; Anell, Thybo & Rasmussen, 2012). The climatic model describes how climatic changes and isostatic adjustments are the controlling factors regarding sediment dispersal and depocentre migration in the eastern North Sea (e.g. Huuse & Clausen, 2001; Huuse, Lykke-Andersen & Michelsen, 2001; Nielsen *et al.* 2009, 2010; Goleadowski, Nielsen & Clausen, 2012). However, more regional studies support the theory that both regional tectonics and climate changes influenced western Scandinavia during Cenozoic time. These studies show that the prograding successions described in this study correlate to a period of substantial lithospheric reorganization in the north Atlantic (e.g. Gaina, Gernigon & Ball, 2009), including alpine compression in southern Europe (e.g. Coward & Dietrich, 1989), and the global cooling event at

the Eocene–Oligocene boundary (e.g. Haq, Hardenbol & Vail, 1987; Abreu & Anderson, 1998; Miller *et al.* 2005; DeConto *et al.* 2008; Pekar & Christie-Blick, 2008; Scher *et al.* 2011). The aim of this paper is to discuss the creation of accommodation space and sediment infill according to the observations presented in the previous section, with emphasis on both regional tectonics and climate changes.

6.a. Creation of accommodation space, climate and shifting depocentres

This section discusses the development and migration of depocentres in relation to well-known eustatic sea-level changes. The base Oligocene unconformity (Fig. 8) corresponds in time with eustatic sea-level fall at the Eocene–Oligocene boundary (e.g. Haq, Hardenbol & Vail, 1987; Miller *et al.* 2005; Van Simaey *et al.* 2005; Fig. 15), concurrent with major glaciation in Antarctica (e.g. Lear *et al.* 2004; Tripathi *et al.* 2005; Pekar & Christie-Blick, 2008). Furthermore, it correlates with the regional O1 unconformity in the eastern North Sea (Huuse & Clausen, 2001; Huuse, 2002; Stoker *et al.* 2005; Japsen *et al.* 2007). A marine origin for the unconformity is suggested from the shape of the erosional surface which resembles marine incision, possibly from ocean currents. Marine incision at the Eocene–Oligocene boundary has also been proposed by Clausen *et al.* (1999) and Huuse & Clausen (2001) for offshore Denmark, to the east of the study area.

The sea-level rise associated with deposition of the TST at the base of OSS-1 is correlative with a rise in eustatic sea level of about 30 m during the early phase of deposition of OSS-1 (Kominz & Pekar, 2001; Miller *et al.* 2005; Fig. 15). However, when considering that water depth was *c.* 100 m during deposition of the RST of the same sequence, additional basin subsidence is required. In particular, one salt structure shows re-activation and diapir growth in the central part of the OSS-1 depocentre. Subsidence from salt removal in the subsurface may have led to additional basin subsidence during early Rupelian time (halokinesis is discussed in more detail in Section 6.b). This is a reflection of how the accommodation space related to OSS-1 developed mainly during marine erosion at the Eocene–Oligocene boundary, combined with halokinesis and rise in eustatic sea level during early Rupelian time.

Succeeding progradation and deposition of the RST of OSS-1 and erosion of the clinofold topsets indicates that the accommodation space created during early Rupelian time was infilled by the RST of the same sequence. The erosional character of the upper bounding surface of OSS-1 (Fig. 7) may reflect either: (1) subaerial exposure and continental incision; (2) erosion and truncation from deep marine currents; or (3) a shallow marine ravinement surface. Both subaerial and deep marine erosion are often recognized by incision, parallel and/or perpendicular to the direction of progradation. Deep marine erosion/incision often occurs basinwards of the clinofold break point, which has




Figure 15. (Colour online) Summary of sea-level changes and regional subsidence during deposition of the individual sequences from this study compared to the eustatic sea-level changes proposed by Kominz & Pekar (2001). Note the vast sediment volume rate in OSS-4.

previously been documented to have occurred during Cenozoic time in the North Sea Basin (Clausen *et al.* 1999; Huuse & Clausen, 2001). The SU at the top of OSS-1 signifies clinoforn tolap truncation (Fig. 7). Incised valleys that may reflect deep marine or continental incision have not been observed at the top of OSS-1 anywhere in the study area. We therefore propose that shallow marine erosion formed the SU at the top of OSS-1, supported by the observation of shallow water fossil assemblages in well 11/10-1, indicating deposition in the middle neritic zone (Fig. 4). Although seismic resolution of the dataset should be adequate to identify channel incision, also observed at the top of OSS-2 (Fig. 12), it is noteworthy that similar

features may be present but are below seismic resolution at the top of sequence OSS-1. However, based on the character of the erosional surface at the top of OSS-1, it is suggested here that the accommodation space available during early Rupelian time was infilled by OSS-1 sediments. As the upper sequence unconformity corresponds to the O1a global cooling event (Pekar *et al.* 2002; Sliwiska & Heilmann-Clausen, 2011), it is however also likely that a drop in eustatic sea level took place at the later phase of deposition of OSS-1.

Following maximum regression and tolap truncation during early Rupelian time (top of OSS-1), aggradation and subsequent flooding of the RST of sequence OSS-1 indicates a rise in relative sea level and the

creation of new accommodation space. The aggrading TST in the lower part of OSS-2 corresponds to a period of increased eustatic sea level (Kominz & Pekar, 2001; Fig. 15). This was followed by a lowering of the eustatic sea level of the same magnitude, which indicates that global sea-level changes alone cannot explain the development of new accommodation space involving an increase in water depth of a minimum of 250 m (Fig. 15). In general, isostatic flexuring due to denudation of the hinterland and sediment loading are important factors regarding subsidence in a sedimentary basin (Watts, Karner & Steckler, 1982; Reynolds, Steckler & Coakley, 1991), and probably also during late Palaeogene and Neogene time in the North Sea (e.g. Huuse, 2002; Goleadowski *et al.* 2012). These processes are considered to be slow, which may explain subsidence patterns over longer periods on the scale of first-order sequences (e.g. the complete Oligocene sedimentary succession in the North Sea; Watts, Karner & Steckler, 1982; Cloetingh *et al.* 1990). Exceptions are when isostatic rejuvenations are related to glacial meltdown where isostatic adjustments of hundreds of metres may take place within a period as short as thousands of years, as revealed during post-glacial rebound in Scandinavia during Quaternary time (e.g. Riis, 1996). The northern hemisphere was most likely devoid of continental glaciers during Oligocene time (e.g. DeConto *et al.* 2008); isostatic rejuvenations therefore fail to explain basin subsidence and the rapid creation of accommodation space over the short time periods of the Oligocene seismic sequences in the eastern North Sea. The creation of accommodation space of *c.* 250 m during late early Oligocene time cannot be solely explained by isostatic flexuring effects caused by increased erosion and sedimentation rates in the hinterland. This is supported by the low volume rates (accumulation rates) during late Rupelian time (Fig. 15), which indicate drier climatic conditions and lower erosion rates in the hinterland. It is therefore proposed here that the accommodation space related to the RST of sequence OSS-2 was created during tectonic uplift of southern Norway and corresponding basin subsidence in the eastern North Sea. The fact that sequence OSS-2 has a uniform thickness across the Fjerritslev Fault Zone area indicates that the northernmost parts of the study area were also likely influenced by basin subsidence. This implies that the sedimentary basin extended northwards, potentially across parts of the present southern Norwegian mainland.

The deposition of OSS-3 and OSS-4 reflects continued creation of accommodation space due to the relative rise of sea level during Chattian time. Lowering of the eustatic sea level took place during deposition of OSS-3 (Fig. 15; Kominz & Pekar, 2001), which may indicate that the creation of accommodation space within the study area had a more local origin. However, the accretionary TST in the lower part of the sequence, combined with the basinwards shift in progradation compared to the RST of OSS-2, support the theory that the creation of accommodation space may have been related to a combination of long-term flexural isostatic effects and

sediment compaction. The accommodation space relative to sequence OSS-3 was completely infilled by the RST of the same sequence, based on clinoform top-set truncation at the upper bounding sequence unconformity (Fig. 13). This is supported by microfossil assemblages, indicating sediment deposition in the inner to middle shoreface in the 11/10–1 well area (Fig. 4). An early Chattian global cooling period (Kominz & Pekar, 2001) may explain the increase in accumulated sediment volumes compared to the Rupelian OSS-1 and OSS-2 sequences, since wet-based alpine glaciers may have developed in the hinterland (DeConto *et al.* 2008; Nielsen *et al.* 2009). This is supported by the SU of OSS-3 corresponding to a period of forced regression in the North Sea area during maximum regression (Clausen, Sliwinska & Goleadowski, 2012). The lowering of the sea level was related to a global cooling event, which appears to have been synchronous with the Oi2 cooling event (Miller *et al.* 1998; Clausen, Sliwinska & Goleadowski, 2012).

Following exposure of the shelf corresponding to the SU at top of OSS-3, new accommodation space with a water depth of *c.* 300 m was created (Fig. 6). The eustatic sea level fluctuated by tens of metres during late Chattian time, although with an overall rise of only *c.* 20 m (Fig. 15; Kominz & Pekar, 2001). This indicates that additional basin subsidence is required to explain the remaining accommodation space available during infill of the RST of OSS-4. The accumulation rate of OSS-4 is three times higher relative to sequence OSS-3 (Fig. 15), implying that the erosion rate during deposition of OSS-4 must have been significantly higher than during early Chattian time (OSS-3). This may be related to the fact that the lower part of sequence OSS-4 corresponds in time with the Oi-2b cooling phase (Van Simaey *et al.* 2005; Sliwinska, Clausen & Heilmann-Clausen, 2010), dated at *c.* 27.1 Ma (Miller *et al.* 1998), increasing the potential of build-up of alpine glaciers in the hinterlands. However, when considering that global cooling also affected the OSS-3 sequence, similar accumulation rates for OSS-3 and OSS-4 would have been expected. The creation of accommodation space related to OSS-4 was rapid, considering the non-accretionary TST at the base of the sequence. Since isostatic rejuvenation related to glacial meltdown is unlikely to have occurred during Oligocene time (DeConto *et al.* 2008), the rapid creation of accommodation space is most likely related to tectonic subsidence. The tectonic subsidence may have taken place in conjunction with tectonic uplift of the hinterland, which may further explain the three-fold increase in accumulation rate during OSS-4 relative to OSS-3.

6.a.1. Mechanisms causing basin subsidence

The exact mechanism responsible for the creation of accommodation space during Oligocene time is still being debated (Faleide *et al.* 2002; Anell, Thybo & Stratford, 2010). However, observations from the current study indicate that basin subsidence is related to




Figure 16. (Colour online) Position of the main depocentres of the OSS-sequences and illustration of drainage pattern during Oligocene time. Uplift of southern Norway may have forced the drainage system eastwards, with the result of sediment transportation mainly from the NNW and north during early Oligocene time and from the northeast during late Oligocene time.

regional tectonic processes. This implies that processes in the hinterland and in the basinal realm acted simultaneously, and that the same mechanism simultaneously created both uplift in the hinterland and subsidence in the eastern North Sea area. No indications of major tectonic displacement of local origin are observed within the basin. This indicates that basin subsidence is related to large-scale lithospheric processes, such as differential vertical movements during compression of the lithosphere (lithospheric folding) or mantle processes. Both processes may cause long-wavelength deformation of the lithosphere of up to hundreds of kilometres, and the depocentres are characterized by accelerated basin subsidence (Vågnes, Gabrielsen & Haremo, 1998; Bourgeois *et al.* 2007; Cloetingh & Burov, 2011) that triggered potential non-accretionary TSTs. However, the exact mechanism is still debated, and the reader is referred to Faleide *et al.* (2002) and Anell, Thybo & Stratford (2010) for an overview of the discussion.

6.a.2. Shifting depocentres and drainage system

Although progradation direction may be influenced by several basinal processes (e.g. wave and tidal currents) with longshore transport, changes in the direction of progradation may reflect changes in the drainage system along with basin subsidence (Fig. 16). However, the gradual shift in progradation from NNW–SSE during earliest Chattian time (OSS-1) to NE–SW during Rupelian time (OSS-3 and OSS-4) may reflect a dynamic fluvial drainage system in the hinterland. The first progradational sequence into the eastern North Sea during early Oligocene time (OSS-1) represents the most severe shift in sediment transportation dur-

ing Cenozoic time. The transportation of only minor amounts of erosional material into the eastern North Sea during Palaeocene–Eocene time indicates that the Oligocene sequences represent sediment deposition related to the initial tectonic uplift phase of southern Norway in late Palaeogene – Neogene time.

The structural high along the Fjerritslev Fault Zone may have acted as a barrier during the deposition of OSS-1 and the lower part of OSS-2, forcing the sediment input to the west with subsequent south-eastwards dispersal (Fig. 15).

It is interpreted here that the drainage system shifted further eastwards during deposition of the upper part of OSS-2 and OSS-3 and OSS-4, allowing fluvial drainage systems to deliver detritus to the basin more from the north and northeast. Since only minor proportions of the Oligocene sedimentary succession in the northern North Sea have been sourced from southern Norway (e.g. Galloway *et al.* 1993; Jordt *et al.* 1995; Jordt, Thyberg & Nøttvedt, 2000; Faleide *et al.* 2002; Anell, Thybo & Artimieva, 2009), the watershed is likely to have been located far to the west, causing most of the erosional material to be transported south and south-eastwards. We believe the uplift of southern Norway has influenced the drainage pattern, as reflected by the shift in sediment transport direction and subsequent direction of progradation. As uplift continued more landmasses were gradually exposed, forcing the drainage system to follow established N–S- and NE–SW-striking lineaments on present-day eastern and southern Norway. This probably continued during Miocene time, when the lower Miocene deposits of onshore Denmark were sourced from southern Norway (Olivarius, 2009) and south-westwards sediment dispersal and depocentre migration continued into the Central Graben area (e.g. Jarsve *et al.* 2014).

6.b. Halokinesis

The prime mechanism for halokinesis in the study area generally appears to be related to differential loading. This often developed in sedimentary basins as prograding successions created greater loading in some parts of the depositional area compared to elsewhere (e.g. Trusheim, 1960; Seni & Jackson, 1983; Hughes & Davison, 1993; Kossow *et al.* 2000; Steward, 2007). However, we cannot exclude the effect of active tectonism on salt mobilization in the study area, which was the prime mechanism regarding salt mobilization onshore Denmark during early Oligocene time (Japsen *et al.* 2007).

The focus of this study is the intrinsic control on sedimentation caused by salt diapirism; one salt diapir in particular influenced the main depocentre of OSS-1. The actual size of the salt diapir appears to be limited in size relative to the size of the depocentre for it to have influenced basin development, however. We therefore believe halokinesis was insignificant with respect to the creation of accommodation space during early Oligocene time (OSS-1). However, halokinesis does seem

to have influenced maximum progradation for OSS-1, OSS-2 and OSS-3 in parts of the depocentres, indicated by the thinning of all the seismic sequences across salt diapirs at the margins of the depocentres. Parts of the main depocentre infilled by OSS-1 are located around a salt structure located WSW of well 11/10–1 (Figs 7, 9), onlapping against the salt structure. These observations show that the salt diapirs form intra-basinal topography, affecting sediment progradation. This effect also influences OSS-2 and OSS-3 progradation (Fig. 9). Sediment progradation is observed on either side of the salt diapirs, supporting the theory that halokinesis had an intrinsic control on sediment progradation. These observations indicate that halokinesis controlled the extension of maximum progradation and exerted an internal control of progradation within the main depocentres.

7. Conclusions

We have shown that renewed interpretation of the Oligocene succession, combining seismic sequence stratigraphic techniques, biostratigraphy and Sr-isotope stratigraphy and published climate data, provides an improved understanding of the coupling of tectonic and climatic influence on the Oligocene source to sink system in the eastern North Sea area.

Each of the four seismic sequences described in this article are characterized as T-R sequences, with a transgressive systems tract (TST) overlain by a regressive systems tract (RST). In OSS-1 and OSS-4 the TST are non-accretionary, while in OSS-2 and OSS-3 they are aggradational. Accommodation space developed simultaneously with initial sediment progradation from NNW. The main findings from this study are as follows.

- New age constraints from this study show that the sediment progradation from southern Norway was initiated during earliest Oligocene time and slightly post-dates the global cooling event at the Eocene–Oligocene boundary.
- It is documented here that the vast increases in water depths during deposition of the Oligocene seismic sequences were related to basin subsidence that outpaced a generally shallowing fall in eustatic sea level. It has also been shown that the creation of accommodation spaces was out of phase relative to intra-Oligocene eustatic sea-level fluctuations.
- Basin subsidence is likely to have accompanied tectonic rise of the southern Norwegian landmass, where the eastern part of the North Sea basin experienced subsidence forced by differential vertical movements of the lithosphere. These vertical movements gave rise to rapid changes in palaeowater depth and accompanying changes from potential subaerial exposure of basin platforms to marine flooding and water depths of up to *c.* 300 m.
- The shift in progradation direction from southeast during earliest Rupelian time to south and southwest

during late Rupelian – Chattian time was a response to tectonic uplift in the western part of southern Norway, which may have forced the drainage system eastwards.

- Within the basin, halokinesis influenced the lateral extension of sediment progradation throughout Rupelian – early Chattian time. The halokinesis was caused by differential sediment loading in adjacent areas of the depositional basin. However, the limited sizes of the salt diapirs means that they are not considered to have influenced basin subsidence.

We have demonstrated that the creation and infill of accommodation spaces during Oligocene time in the eastern North Sea are related to both tectonic processes and climate changes. This study also highlights the importance of studying the interplay between tectonics and climate in order to better understand the source to sink relationship.

Acknowledgements. The authors would like to thank Fugro and TGS Nopec for making the seismic data available for this study. Erik S. Rasmussen, Mette Danielsen, Mads Huuse, Peter Japse and Amanadine Prelat are thanked for insightful and constructive comments on this paper. We are also grateful to Adrian Read for proofreading and commenting on the manuscript. We would also like to thank Rune Goa of NPD for improving the figures in the paper.

References

- ABREU, V. S. & ANDERSON, J. B. 1998. Glacial eustacy during the Cenozoic: sequence stratigraphic implications. *AAPG Bulletin* **82**, 1385–400.
- ANELL, I., THYBO, H. & ARTIMEVA, I. M. 2009. Cenozoic uplift and subsidence in the North Atlantic Region. *Tectonophysics* **474**, 78–105.
- ANELL, I., THYBO, H. & RASMUSSEN, E. 2012. A synthesis of Cenozoic sedimentation in the North Sea. *Basin Research* **23**, 1–26.
- ANELL, I., THYBO, H. & STRATFORD, W. 2010. Relating Cenozoic North Sea sediments to topography in southern Norway: the interplay between tectonics and climate. *Earth and Planetary Science Letters* **300**, 19–32.
- BERGGREN, W. A., KENT, D. V., SWISHER, C. C. & AUBRY, M.-P. 1995. A revised Cenozoic geochronology and chronostratigraphy. In *Geochronology Time Scale and Global Stratigraphic Correlation* (ed. W. A. Berggren), pp. 129–212. Society for Sedimentary Geology, Special Publication no. 54.
- BERGGREN, W. A., KENT, D. V. & VAN COUVERING, J. A. 1985. The Neogene. Part 2. Neogene geochronology and chronostratigraphy. In *The Chronology of the Geological Record* (ed. N. J. Snelling), pp. 141–95. Geological Society of London, Memoir no. 10.
- BERTHELSEN, A. 1998. The Tornquist Zone northwest of the Carpathians: an intraplate pseudosuture. *Geologiska Foreningen* **120**, 223–30.
- BULLIMORE, S., HENRIKSEN, S., LIESTØL, F. M. & HELLAND-HANSEN, W. 2005. Clinoform stacking patterns, shelf-edge trajectories and facies associations in Tertiary coastal deltas, offshore Norway: implications for the prediction of lithology in prograding systems. *Norwegian Journal of Geology* **85**, 169–87.

- BOURGOIS, O., FORD, M., DIRAISON, M., VESLUD, C. L. C. D., GERBAULT, M., PIK, R., RUBY, N. & BONNET, S. 2007. Separation of rifting and lithospheric folding signatures in the NW-Alpine foreland. *International Journal of Earth Sciences* **96**, 1003–31.
- CANDE, S. C. & KENT, D. V. 1992. A new geomagnetic polarity time-scale for the late Cretaceous and Cenozoic. *Journal of Geophysical Research* **97**, 13917–51.
- CHRISTENSEN, L. & ULLEBERG, K. 1973. Sedimentology and micropalaeontology of the Middle Oligocene sequence at Sofienlund – Denmark. *Geological Society of Denmark, Bulletin* **22**, 283–305.
- CLAUSEN, O. R., GREGERSEN, U., MICHELSEN, O. & SØRENSEN, J. C. 1999. Factors controlling the Cenozoic sequence development in the eastern parts of the North Sea. *Journal of Geological Society* **156**, 809–16.
- CLAUSEN, O. R., SLIWINSKA, K. K. & GOLEDOWSKI, B. 2012. Oligocene climate changes controlling forced regression in the eastern North Sea. *Marine and Petroleum Geology* **29**, 1–14.
- CLOETINGH, S. & BUROV, E. 2011. Lithospheric folding and sedimentary basin evolution: a review and analysis of formation mechanisms. *Basin Research* **23**, 257–90.
- CLOETINGH, S., GRADSTEIN, F. M., KOOL, H., GRANT, A. C. & KAMINSKI, M. 1990. Plate reorganization: a cause of rapid late Neogene subsidence and sedimentation around the North Atlantic. *Journal of the Geological Society* **147**, 495–506.
- COLEMAN, J. M. 1988. Dynamic changes and processes in the Mississippi River delta. *Geological Society of America Bulletin* **100**, 999–1015.
- COWARD, M. & DIETRICH, D. 1989. Alpine tectonics – an overview. In *Alpine Tectonics* (eds M. P. Coward, D. Dietrich and R. G. Park), pp. 1–29. Geological Society of London, Special Publication no. 45.
- DANIELSEN, M., MICHELSEN, O. & CLAUSEN, O. R. 1997. Oligocene stratigraphy and basin development in the Danish North Sea sector based on log interpretations. *Marine and Petroleum Geology* **14**, 931–51.
- DECONTO, R. M., POLLARD, D., WILSON, P. A., PALIKE, H., LEAR, C. H. & PAGANI, M. 2008. Thresholds for Cenozoic bipolar glaciation. *Nature* **455**, 652–6.
- DMITRIEVA, E., JACKSON, C. A.-L., HUUSE, M. & MCCARTHY, A. 2012. Paleocene deep-water depositional systems in the North Sea Basin - a 3D seismic and well data case study, offshore Norway. *Petroleum Geoscience* **18**, 97–114.
- EIDVIN, T., RIIS, F., RASMUSSEN, E. S. & RUNDBERG, Y. 2013. Investigation of Oligocene to Lower Pliocene deposits in the Nordic offshore area and onshore Denmark. *NPD Bulletin* **10**, 62.
- EIDVIN, T. & RUNDBERG, Y. 2001. Late Cainozoic stratigraphy of the Tampen area (Snorre and Visund fields) in the northern North Sea, with emphasis on the chronology of early Neogene sands. *Norsk Geologisk Tidsskrift* **81**, 119–60.
- EIDVIN, T. & RUNDBERG, Y. 2007. Post-Eocene strata of the southern Viking Graben, northern North Sea; integrated biostratigraphic, strontium isotopic and lithostratigraphic study. *Norwegian Journal of Geology* **87**, 391–450.
- EMBRY, A. 1988. Triassic sea-level changes: evidence from Canadian Arctic Archipelago. In *Sea-level Changes – An Integrated Approach* (eds C. K. Wilgus, B. S. Hastings, H. Posamentier, J. V. Wagoner, C. A. Ross and C. Kendall), pp. 249–59. SEPM, Special Publication no. 42.
- EMBRY, A. 1993. Transgressive-regressive (T-R) sequence analysis of the Jurassic succession of the Sverdrup Basin, Canadian Arctic Archipelago. *Canadian Journal of Earth Sciences* **30**, 301–20.
- EMBRY, A. 1995. Sequence boundaries and sequence hierarchies: problems and proposals. In *Sequence Stratigraphy on the Northwest European Margin* (eds R. Steel, V. L. Felt, E. P. Johannessen and C. Mathieu), pp. 1–11. Norwegian Petroleum Society, Special Publication no. 5.
- EMBRY, A. & JOHANNESSEN, E. 1992. T-R sequence stratigraphy, facies analysis reservoir distribution in the uppermost Triassic–Lower Jurassic succession, western Sverdrup Basin, Arctic Canada. In *Arctic Geology and Petroleum Potential* (eds T. Vorren, E. Bergsager, Ø. A. Dahl-Stamnes, E. Holter, B. Johansen, E. Lie and T. B. Lund), pp. 121–46. Norwegian Petroleum Society, Special Publication no. 2.
- FALEIDE, J. I., KYRKJEBØ, R., KJENNERUD, T., GABRIELSEN, R. H., JORDT, H., FANAVOLL, S. & BJERKE, M. D. 2002. Tectonic impact on Sedimentary processes during Cenozoic evolution of the Northern North Sea and surrounding areas. In *Exhumation of the North Atlantic Margin: Timing, Mechanisms and Implications for Petroleum Exploration* (eds A. G. Dorè, J. A. Cartwright, M. S. Stoker, J. P. Turner & N. White), pp. 235–69. Geological Society of London, Special Publication no. 196.
- FREDRIKSEN, S., NIELSEN, S. B. & BALLING, N. 2001. A numerical dynamic model for the Norwegian-Danish Basin. *Tectonophysics* **343**, 165–83.
- GABRIELSEN, R. H., BRAATHEN, A., OLESEN, O., FALEIDE, J. I., KYRKJEBØ, R. & REDFIELD, T. F. 2005. Vertical movements of the South-Western Fennoscandia: a discussion of regions and processes from the present to the Devonian. In *Onshore-Offshore Relationships on the North Atlantic Margin, Proceedings of the Norwegian Petroleum Society Conference* (eds B. T. G. Wandås, J. P. Nystuen, E. Eide and F. Gradstein), pp. 1–28. Norwegian Petroleum Society, Special Publication no. 12.
- GABRIELSEN, R. H., FALEIDE, J. I., PASCAL, C., BRAATHEN, A., NYSTUEN, J. P., ETZELMULLER, B. & O'DONNELL, S. 2010. Latest Caledonian to present tectonomorphological development of Southern Norway. *Marine and Petroleum Geology* **27**, 709–23.
- GAINA, C., GERNIGON, L. & BALL, P. 2009. Palaeocene–Recent plate boundaries in the NE Atlantic and the formation of the Jan Mayen microcontinent. *Journal of the Geological Society of London* **166**, 601–16.
- GALLOWAY, W. E., GARBER, J. L., LIU, X. & SLOAN, B. J. 1993. Sequence stratigraphic and depositional framework of the Cenozoic fill, Central and Northern North Sea Basin. In *Petroleum Geology of Northwest Europe: Proceedings of the 4th Conference* (ed. J. R. Parker), pp. 33–43. Geological Society of London, Petroleum Geology Conference Series no. 4.
- GOLEDOWSKI, B., NIELSEN, S. B. & CLAUSEN, O. R. 2012. Patterns of Cenozoic sediment flux from western Scandinavia. *Basin Research* **23**, 1–24.
- GRADSTEIN, F. & BÄCKSTRÖM, S. 1996. Cainozoic biostratigraphy and paleobathymetry, northern North Sea and Haltenbanken. *Norsk Geologisk Tidsskrift* **76**, 3–32.
- HAMBERG, L., DAM, G., WILHELMSON, C. & OTTESEN, T. G. 2005. Paleocene deep-marine sandstone plays in the Siri Canyon, offshore Denmark – southern Norway. In *Petroleum Geology: North-West Europe and Global Perspectives* (eds A. G. Dorè and B. A. Vining), pp. 1185–98. Geological Society of London, Petroleum Geology Conference Series no. 6.
- HANSEN, J. P. V. & RASMUSSEN, E. S. 2008. Structural, sedimentologic, and sea-level controls on sand

- distribution in a steep-clinoform asymmetric wave-influenced delta: Miocene Billund Sand, Eastern Danish North Sea and Jylland. *Journal of Sedimentary Research* **78**, 130–46.
- HAO, B. U., HARDENBOL, J. & VAIL, P. R. 1987. Chronology of fluctuating sea levels since the Triassic. *Science* **235**, 1156–67.
- HEEREMANS, M. & FALEIDE, J. I. 2004. Late Carboniferous–Permian tectonics and magmatic activity in the Skagerrak, Kattegat and the North Sea. In *Permo-Carboniferous Magmatism and Rifting in Europe* (eds M. Wilson, E.-R. Neumann, G. R. Davies, M. J. Timmerman, M. Heeremans & B. T. Larsen), pp. 157–76. Geological Society of London, Special Publication no. 223.
- HEEREMANS, M., TIMMERMAN, M. J., KIRSTEIN, L.A. & FALEIDE, J. I. 2004. New constraints on the timing of late Carboniferous–early Permian volcanism in the central North Sea. In *Permo-Carboniferous Magmatism and Rifting in Europe* (eds M. Wilson, E.-R. Neumann, G. R. Davies, M. J. Timmerman, M. Heeremans & B. T. Larsen), pp. 177–93. Geological Society of London, Special Publication no. 223.
- HELLAND-HANSEN, W. 1995. Sequence stratigraphy theory: remarks and recommendations. In *Sequence Stratigraphy on the Northwest European Margin* (eds R. J. Steel, V. L. Felt, E. P. Johannessen and C. Mathieu), pp. 13–21. Norwegian Petroleum Society, Special Publication no. 5.
- HOWARTH, R. J. & MCARTHUR, J. M. 1997. Statistics for strontium isotope stratigraphy: A robust LOWESS fit to marine Sr-isotope curve for 0 to 206 Ma, with look-up table for derivation of numeric age. *Journal of Geology* **105**, 441–56.
- HUGHES, M. & DAVISON, I. 1993. Geometry and growth kinematics of salt pillows in the southern North Sea. *Tectonophysics* **228**, 239–54.
- HUUSE, M. 2002. Cenozoic uplift and denudation of southern Norway: insights from the North Sea Basin. In *Exhumation of the North Atlantic Margin: Timing, Mechanisms and Implications for Petroleum Exploration* (eds A. G. Dorè, J. A. Cartwright, M. S. Stoker, J. R. Turner & N. White), pp. 209–33. Geological Society of London, Special Publication no. 196.
- HUUSE, M. & CLAUSEN, O. R. Clausen 2001. Morphology and origin of major Cenozoic sequence boundaries in the eastern North Sea basin: top Eocene, near-top Oligocene and the mid-Miocene unconformity. *Basin Research* **13**, 17–41.
- HUUSE, M., LYKKE-ANDERSEN, H. & MICHELSEN, O. 2001. Cenozoic evolution of the eastern Danish North Sea. *Marine Geology* **177**, 243–69.
- JAPSEN, P., BIDSTRUP, T. & LIDMAR-BERGSTRÖM, K. 2002. Neogene uplift and erosion of southern Scandinavia induced by the rise of the South Swedish Dome. *Geological Society, London* **196**, 183–207.
- JAPSEN, P., BONOW, J. M., GREEN, P. F., CHALMERS, J. A. & LIDMAR-BERGSTRÖM, K. 2006. Elevated, passive continental margins: Long-term highs or Neogene uplifts? New evidence from West Greenland. *Earth and Planetary Science Letters* **248**, 330–9.
- JAPSEN, P. & CHALMERS, J. A. 2000. Neogene uplift and tectonics around the North Atlantic: overview. *Global and Planetary Change* **24**, 165–73.
- JAPSEN, P., GREEN, P. F., NIELSEN, L. H., RASMUSSEN, E. S. & BIDSTRUP, T. 2007. Mesozoic–Cenozoic exhumation events in the eastern North Sea Basin: a multi-disciplinary study based on palaeothermal, palaeoburial, stratigraphic and seismic data. *Basin Research* **19**, 451–90.
- JARSVE, E. M., FALEIDE, J. I., GABRIELSEN, R. H. & NYSTUEN, J. P. 2014. Mesozoic and Cenozoic basin configurations in the North Sea. In *From Depositional Systems to Sedimentary Successions on the Norwegian Continental Margin* (eds A. W. Martinius, R. Ravnås, J. Howell, T. Olsen, R. J. Steel and J. Wonham), pp. 417–52. International Association of Sedimentologists, Special Publication no. 46.
- JORDT, H., FALEIDE, J. I., BJØRLYKKE, K. & IBRAHIM, M. T. 1995. Cenozoic sequence stratigraphy of the central and northern North Sea Basin: tectonic development, sediment distribution and provenance areas. *Marine and Petroleum Geology* **12**, 845–79.
- JORDT, H., THYBERG, B. I. & NØTTVEDT, A. 2000. Cenozoic evolution of the central and northern North Sea with focus on differential vertical movements of the basin floor and surrounding clastic source areas. In *Dynamics of the Norwegian Margin* (eds A. Nøttvedt, B. T. Larsen, R. H. Gabrielsen, S. Olaussen, H. Brekke, B. Torudbakken, O. Birkelund & J. Skogseid), pp. 219–43. Geological Society of London, Special Publication no. 167.
- KING, C. 1989. Cenozoic of the North Sea. In *Stratigraphical Atlas of Fossils Foraminifera* (ed. D. G. Jenkins & J. W. Murray), pp. 418–89. Chichester: Ellis Horwood Ltd.
- KOMINZ, M. A. & PEKAR, S. F. 2001. Oligocene eustasy from two-dimensional sequence stratigraphic backstripping. *Geological Society of America Bulletin* **113**, 291–304.
- KOSSOW, D., KRAWCZYK, C., MCCANN, T., STRECKER, M. & NEGENDANK, J. F. W. 2000. Style and evolution of salt pillows and related structures in the northern part of the Northeast German Basin. *International Journal of Earth Sciences* **89**, 652–64.
- LEAR, C. H., ROSENTHAL, Y., COXALL, H. K. & WILSON, P. A. 2004. Late Eocene to early Miocene ice sheets dynamics and the global carbon cycle. *Paleoceanography* **19**, 1–11.
- MARTINSEN, O. J., BOEN, F., CHARNOCK, M. A., MANGERUT, G. & NØTTVEDT, A. 1999. Cenozoic development of the Norwegian margin 60–64°N: sequences and sedimentary response to variable basin physiography and tectonic setting. In *Petroleum Geology of NW Europe, Proceedings of the 5th Conference* (eds A. J. Fleet and S. A. R. Boldy), pp. 293–304. Geological Society of London, Petroleum Geology Conference Series no. 5.
- MCARTHUR, J. M., HOWARTH, R. J. & BAILEY, T. R. 2001. Strontium isotope stratigraphy: LOWESS version 3: best fit to the marine Sr-isotope curve for 0–509 Ma and accompanying look-up table for deriving numerical age. *Journal of Geology* **109**, 155–70.
- MEDVEDEV, S., SOUCHE, A. & HARTZ, E. H. 2013. Influence of ice sheet and glacial erosion on passive margins of Greenland. *Geomorphology* **193**, 36–46.
- MICHELSEN, O. & DANIELSEN, M. 1996. Sequence and systems tract interpretation of the epicontinental Oligocene deposits in the Danish North Sea. In *Geology of Siliciclastic Shelf Seas* (eds M. De Batist & P. Jacobsen), pp. 1–13. Geological Society of London, Special Publication no. 117.
- MICHELSEN, O., DANIELSEN, M., HEILMANN-CLAUSEN, C., JORDT, H., LAURSEN, G. V. & THOMSEN, E. 1995. Occurrence of major sequence stratigraphic boundaries in relation to basin development in the Cenozoic deposits of the southeastern North Sea. In *Sequence Stratigraphy of the Northwestern European Margin* (ed. R. J. Steel),

- pp. 415–27. Norwegian Petroleum Society (NPF), Special Publication no. 5.
- MICHELSSEN, O., THOMSEN, E., DANIELSEN, M., HEILMANN-CLAUSEN, C., JORDT, H. & LAURSEN, G. V. 1998. Cenozoic sequence stratigraphy in the eastern North Sea. In *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins* (eds P.Ch. De Graciansky, T. Jacquin and P. R. Vail), pp. 91–118. Society for Sedimentary Research, Special Publication no. 60.
- MILLER, K. G., KOMINZ, M. A., BROWNING, J. V., WRIGHT, J. D., MOUNTAIN, G. S., KATZ, M. E., SUGARMAN, P. J., CRAMER, B. S., CHRISTIE-BLICK, N. & PEKAR, S. F. 2005. The phanerozoic record of global sea-level change. *Science* **310**, 1293–8.
- MILLER, K. G., MOUNTAIN, G. S., BROWNING, J. V., KOMINZ, M., SUGARMAN, P. J., CHRISTIE-BLICK, N., KATZ, M. E. & WRIGHT, J. D. 1998. Cenozoic global sea level, sequences, and the New Jersey transect: results from coastal plain and continental slope drilling. *Reviews of Geophysics* **36**, 569–601.
- MILLER, S. R., SAK, P. B., KIRBY, E. & BIERMAN, P. R. 2013. Neogene rejuvenation of central Appalachian topography: evidence for differential rock uplift from stream profiles and erosion rates. *Earth and Planetary Science Letters* **369–70**, 1–12.
- MITCHUM, R. M. J., VAIL, P. R. & THOMPSON, S. I. 1977. Seismic stratigraphy and global changes in sea-level, part 2: the depositional sequence as the basic unit for stratigraphic analysis. In *Seismic Stratigraphy: Application to Hydrocarbon Exploration* (ed. C. Payton), pp. 53–92. American Association of Petroleum Geologists, Memoir no. 26.
- NICHOLS, G. 2009. *Sedimentology and Stratigraphy*, second edition. West Sussex, UK: Wiley-Blackwell, 419 pp.
- NIELSEN, S. B., CLAUSEN, O. R., JACOBSEN, B. H., THOMSEN, E., HUUSE, M., GALLAGHER, K., BALLING, N. & EGHOLM, D. 2010. The ICE hypothesis stands: how the dogma of late Cenozoic tectonic uplift can no longer be sustained in the light of data and physical laws. *Journal of Geodynamics* **50**, 102–11.
- NIELSEN, S. B., GALLAGHER, K., LEIGHTON, C., BALLING, N., SVENNINGSEN, L., JACOBSEN, B. H., THOMSEN, E., NIELSEN, O. B., HEILMANN-CLAUSEN, C., EGHOLM, D. L., SUMMERFIELD, M. A., CLAUSEN, O. R., PIOTROWSKI, J. A., THORSEN, M. R., HUUSE, M., ABRAHAMSEN, N., KING, C. & LYKKE-ANDERSEN, H. 2009. The evolution of western Scandinavian topography: a review of Neogene uplift versus the ICE (isostasy-climate-erosion) hypothesis. *Journal of Geodynamics* **47**, 72–95.
- OLIVARIUS, M. 2009. Provenance and facies of Miocene sand succession in western Denmark based on bulk geochemistry, heavyminerals and zircon dating. Master's thesis. Department of Geography and Geology, University of Copenhagen. Published thesis.
- PEKAR, S. F. & CHRISTIE-BLICK, N. 2008. Resolving apparent conflicts between oceanographic and Antarctic climate records and evidence for a decrease in pCO₂ during the Oligocene through early Miocene (34–16 Ma). *Palaeogeography, Palaeoclimatology, Palaeoecology* **260**, 41–9.
- PEKAR, S. F., CHRISTIE-BLICK, N., KOMINZ, M. A. & MILLER, K. G. 2002. Calibration between eustatic estimates from backstripping and oxygen isotopic records for the Oligocene. *Geology* **30**, 903–6.
- POSAMENTIER, H. W., JERVEY, M. T. & VAIL, P. R. 1988. Eustatic controls on clastic deposition. I. Conceptual framework. In *Sea Level Changes—An Integrated Approach* (eds C. K. Wilgus, B. S. Hastings, C. G. St. C. Kendall, H. W. Posamentier, C. A. Ross and Van Wagoner), pp. 110–124. Society for Sedimentary Research, Special Publication no. 42.
- POSAMENTIER, H. W. & VAIL, P. R. 1988. Eustatic controls on clastic deposition. II. Sequence and systems tract models. In *Sea Level Changes—An Integrated Approach* (eds C. K. Wilgus, B. S. Hastings, C. G. St. C. Kendall, H. W. Posamentier, C. A. Ross and J. C. Van Wagoner), pp. 125–154. Society for Sedimentary Research, Special Publication no. 42.
- RASMUSSEN, E. S. 2009. Neogene inversion of the Central Graben and Ringkøbing-Fyn High, Denmark. *Tectonophysics* **465**, 84–97.
- REDFIELD, T. F. & OSMUNDSEN, P. T. 2013. The long-term topographic response of a continent adjacent to a hyper-extended margin: a case study from Scandinavia. *Geological Society of America Bulletin* **125**, 184–200.
- REYNOLDS, D. J., STECKLER, M. S. & COAKLEY, B. J. 1991. The role of the sediment load in sequence stratigraphy: the influence of flexural isostasy and compaction. *Journal of Geophysical Research* **96**, 6931–49.
- RIIS, F. 1996. Quantification of Cenozoic vertical movements of Scandinavia by correlation of morphological surfaces with offshore data. *Global and Planetary Change* **12**, 331–57.
- ROHRMAN, M., VAN DER BEEK, P., ANDRIESSEN, P. & CLOETINGH, S. 1995. Meso-Cenozoic morphotectonic evolution of southern Norway: neogene domal uplift inferred from apatite fission track thermochronology. *Tectonics* **14**, 704–18.
- SCHER, H. D., BOHATY, S. M., ZACHOS, J. C. & DELANEY, M. L. 2011. Two-stepping into icehouse: East Antarctic weathering during progressive ice-sheet expansion at the Eocene-Oligocene transition. *Geology* **39**, 383–6.
- SCHJØLER, P., ANDSBJERG, J., CLAUSEN, O. R., DAM, G., DYBKJÆR, K., HAMBER, L., HEILMANN-CLAUSEN, C., JOHANNESSEN, E. P., KRISTENSEN, L. E., PRINCE, & I. Rasmussen, J. A. Rasmussen (eds) 2007. *Lithostratigraphy of the Palaeogene – Lower Neogene Succession of the Danish North Sea*. Geological Survey of Denmark and Greenland, Bulletin no. 12.
- SENI, S. J. & JACKSON, M. P. A. 1983. Evolution of salt structures, East Texas Diapir Province, Part 1: sedimentary record of Halokinesis. *The American Association of Petroleum Geologists Bulletin* **67**, 1219–44.
- SLIWINSKA, K. K., CLAUSEN, O. R. & HEILMANN-CLAUSEN, C. 2010. A mid-Oligocene cooling (Oi-2b) reflected in the dinoflagellate record and in depositional sequence architecture. An integrated study from the eastern North Sea Basin. *Marine and Petroleum Geology* **27**, 1424–30.
- SLIWINSKA, K. K. & HEILMANN-CLAUSEN, C. 2011. Early Oligocene cooling reflected by the dinoflagellate cyst *Svalbardella cooksoniae*. *Palaeogeography, Palaeoclimatology, Palaeoecology* **305**, 138–49.
- STEWART, S. A. 2007. Salt tectonics in the North Sea Basin: a structural style template for seismic interpreters. In *Deformation of the Continental Crust: The Legacy of Mike Coward* (eds A. C. Ries, R. W. H. Butler and R. H. Graham), pp. 361–96. Geological Society of London, Special Publication no. 272.
- STOKER, M. S., PRAEG, D., SHANNON, P. M., HJELSTUEN, B. O., LABER, J. S., NIELSEN, T., VAN WEERING, T. C. E., SEJRUP, H. P. & EVANS, D. 2005. Neogene evolution of the Atlantic continental margin of NW Europe (Lofoten Islands to SW Ireland): anything but passive. In *North-West Europe and Global Perspectives* (eds A. G. Doré & B. A. Vinning), pp. 1057–76. Geological

- society, London, Petroleum Geology Conference Series 6.
- TRIPATI, A., BACKMAN, J., ELDERFIELD, H. & FERRETTI, P. 2005. Eocene bipolar glaciation associated with global carbon cycle changes. *Nature* **436**, 341–6.
- TRUSHEIM, F. 1960. Mechanism of salt migration in Northern Germany. *Bulletin of the American Association of Petroleum Geologists* **44**, 1519–40.
- VÅGNES, E., GABRIELSEN, R. H. & HAREMO, P. 1998. Late Cretaceous – Cenozoic intraplate contractional deformation at the Norwegian continental shelf: timing, magnitude and regional implications. *Tectonophysics* **300**, 29–46.
- VAIL, P. R. 1987. Seismic stratigraphy interpretation procedure. In *Atlas of Seismic Stratigraphy* (ed. A. W. Bally), pp. 1–10. American Association of Petroleum Geologists, Studies in Geology no. 27.
- VAN SIMAEYS, S., BRINKHUIS, H., PROSS, J., WILLIAMS, G. L. & ZACHOS, J. C. 2005. Arctic dinoflagellate migrations mark the strongest Oligocene glaciations. *Geology* **33**, 709–12.
- VAN WAGONER, J. C., MITCHUM JR., R. M., CAMPION, K. M. & RAHMANIAN, V. D. (eds) 1990. *Siliciclastic Sequence Stratigraphy in Well Logs, Core, and Outcrops: Concepts for High-Resolution Correlation of Time and Facies*. American Association of Petroleum Geologists, Methods in Exploration Series 7, 55 pp.
- VAN WAGONER, J. C., MITCHUM, R. M., POSAMENTIER, H. W. & VAIL, P. R. 1987. An overview of sequence stratigraphy and key definitions. In *Seismic Stratigraphy, Volume 1. Studies in Geology* (ed. A. W. Bally), pp. 11–14. American Association of Petroleum Geologists, Studies in Geology no. 27.
- VAN WAGONER, J. C., POSAMENTIER, H. W., MITCHUM, R. M. J., VAIL, P. R., SARG, J. F., LOUTIT, T. S. & HARDENBOL, J. 1988. An overview of the fundamentals of sequence stratigraphy and key definitions. In *Sea-Level Changes—An Integrated Approach* (eds C. K. Wilgus, B. S. Hastings, C. G. S. C. Kendall, H. W. Posamentier, C. A. Ross and J. C. Van Wagoner), pp. 39–45. Society for Sedimentary Geology, Special Publication no. 42.
- VEEKEN, P. C. H. 2007. *Seismic Stratigraphy, Basin Analysis and Reservoir Characterization*. Amsterdam: Elsevier, 509 pp.
- WATTS, A. B., KARNER, G. D. & STECKLER, M. S. 1982. Lithospheric flexure and the Evolution of Sedimentary Basins. *Philosophical Transactions of the Royal Society of London* **305**, 249–81.
- ZIEGLER, P. A. 1990. *Geological Atlas of Western and Central Europe*. Amsterdam: Elsevier, 130 pp.