


Thickness of the Lower Miocene Skade Formation with the extent of the Hutton sand and Lower Miocene mudstones



OD 1209006

Fig. 10: Distribution of Upper Oligocene/Lower Miocene sediments in the northern North Sea with a thickness map of the Skade Formation according to NPD (2011), outline of the Miocene-Upper Pliocene Hutton sand according to Gregersen & Johannessen (2007) and outline of Lower Miocene mudstones according to Rundberg & Eidvin (2005). After Eidvin et al. (2013a).

References

Eidvin, T., Riis, F. & Gjeldvik, I. T., 2013a: The Lower Miocene Skade Formation in the northern North Sea (Extent and thickness, age from fossil and Sr isotope correlations, lithology, paleobathymetry and regional correlation). NGF Abstracts and proceedings, no. 1, 2013, pp. 28-29. Poster available from the internet:

<http://www.npd.no/Global/Norsk/3-Publikasjoner/Presentasjoner/NGF-Vinterkonferanse-2013/Poster-1-til-NGF-vintermotet-nett.pdf>

Gregersen, U. & Johannessen, P. N., 2007: Distribution of the Neogene Utsira Sand and Hutton Sand, and the succeeding deposits in the Viking Graben area, North Sea. *Marine and Petroleum Geology* 24, 591-606.

NPD, 2011: CO2 Atlas, Norwegian North Sea. Available from the internet:

<http://www.npd.no/Global/Norsk/3-Publikasjoner/Rapporter/PDF/CO2-ATLAS-lav.pdf>

Rundberg, Y. & Eidvin, T., 2005: Controls on depositional history and architecture of the Oligocene-Miocene succession, northern North Sea Basin. In B.T.G. Wandaas et al. (eds.): *Onshore-Offshore Relationships on the North Atlantic Margin*. NPF Special Publication 12, 207-239.